

# A $q$ -analogue of perfect matroid designs

Relinde Jurrius

Université de Neuchâtel, Switzerland

Finite fields and applications (Fq13)

June 8, 2017

**Matroid:** a pair  $(E, r)$  with

- ▶  $E$  finite set;
- ▶  $r : 2^E \rightarrow \mathbb{N}_0$  a function, the *rank function*, with for all  $A, B \in E$ :
  - (r1)  $0 \leq r(A) \leq |A|$
  - (r2) If  $A \subseteq B$  then  $r(A) \leq r(B)$ .
  - (r3)  $r(A \cup B) + r(A \cap B) \leq r(A) + r(B)$  (semimodular)

Examples:

- ▶ Set of vectors; rank = matrix rank  
In particular: columns of generator matrix of linear code
- ▶ Set of edges of a graph; rank = size of spanning tree

A subset  $F \subseteq E$  is a **flat** if  $r(F \cup \{x\}) > r(F)$  for any  $x \notin F$ .

The **closure** of a subset  $A \subseteq E$  is the smallest flat that contains  $A$ .

Flats are equal to their closure: *closed sets*.

A matroid is also a pair  $(E, \mathcal{F})$  with

- ▶  $E$  finite set;
- ▶  $\mathcal{F} \subseteq 2^E$  a collection of subsets, the *flats*, with:
  - (F1)  $E \in \mathcal{F}$
  - (F2) If  $F_1, F_2 \in \mathcal{F}$  then  $F_1 \cap F_2 \in \mathcal{F}$ .
  - (F3) If  $F \in \mathcal{F}$ , then every  $x \notin F$  is in a unique flat covering  $F$ .

Theorem (Birkhoff, 1935)

*The poset of flats of a matroid is a co-atomic lattice.*

A **perfect matroid design** is a matroid such that all flats of the same rank have the same size.

### Example

- ▶ (Truncations of) projective spaces;
- ▶ (Truncations of) affine spaces;
- ▶ Steiner systems;
- ▶ Rank 4 PMDs coming from Moufang loops.

Theorem (Murty, Young & Edmonds, 1970)

*The independent sets / circuits / flats of size  $j$  form a design.*

$q$ -analogue: finite set  $\longrightarrow$  finite vector space over  $\mathbb{F}_q$

Example

$$\binom{n}{k} = \text{number of sets of size } k \text{ contained in set of size } n$$

$$\begin{aligned} \left[ \begin{matrix} n \\ k \end{matrix} \right]_q &= \text{number of } k\text{-dim subspaces of } n\text{-dim vector space over } \mathbb{F}_q \\ &= \prod_{i=0}^{k-1} \frac{q^n - q^i}{q^k - q^i} \end{aligned}$$

## Example

$t$ - $(v, k, \lambda)$  **design**: pair  $(X, \mathcal{B})$  with

- ▶  $X$  set with  $v$  elements (points)
- ▶  $\mathcal{B}$  family of subsets of  $X$  of size  $k$  (blocks)
- ▶ Every  $t$ -tuple of points is contained in exactly  $\lambda$  blocks

$t$ - $(v, k, \lambda; q)$  **subspace design**: pair  $(X, \mathcal{B})$  with

- ▶  $X$   $v$ -dim vectorspace over  $\mathbb{F}_q$
- ▶  $\mathcal{B}$  family of  $k$ -dim subspaces of  $X$  (blocks)
- ▶ Every  $t$ -dim subspace is contained in exactly  $\lambda$  blocks

If  $\lambda = 1$  we call the (subspace) design a  $(q)$ -Steiner system

finite set	finite space $\mathbb{F}_q^n$
element	1-dim subspace
size	dimension
$n$	$\frac{q^n - 1}{q - 1}$
intersection	intersection
union	sum
complement	(it depends)

From  $q$ -analogue to 'normal': let  $q \rightarrow 1$ .



**q-Matroid:** a pair  $(E, r)$  with

- ▶  $E$  finite dimensional vector space;
- ▶  $r : \{\text{subspaces of } E\} \rightarrow \mathbb{N}_0$  a function, the *rank function*, with for all  $A, B \subseteq E$ :
  - (r1)  $0 \leq r(A) \leq \dim A$
  - (r2) If  $A \subseteq B$  then  $r(A) \leq r(B)$ .
  - (r3)  $r(A + B) + r(A \cap B) \leq r(A) + r(B)$  (semimodular)

Theorem (J. & Pellikaan, 2016)

*Every  $\mathbb{F}_{q^m}$ -linear rank metric code gives a  $q$ -matroid.*

Proof.

Let  $E = \mathbb{F}_q^n$  and  $G$  be a generator matrix of the code.

Let  $A \subseteq E$  and  $Y$  a matrix whose columns span  $A$ .

$$\boxed{G} \quad \boxed{Y} = \boxed{GY}$$

Then  $r(A) = \text{rk}(GY)$  satisfies the axioms (r1),(r2),(r3). □

A  $q$ -matroid is also a pair  $(E, \mathcal{F})$  with

- ▶  $E$  finite set;
- ▶  $\mathcal{F}$  a collection of subspaces, the *flats*, with:
  - (F1)  $E \in \mathcal{F}$
  - (F2) If  $F_1, F_2 \in \mathcal{F}$  then  $F_1 \cap F_2 \in \mathcal{F}$ .
  - (F3) If  $F \in \mathcal{F}$ , then every 1-dimensional subspace  $x \not\subseteq F$  is in a unique flat covering  $F$ .

Theorem (Crapo, 1964)

*The poset of flats of a  $q$ -matroid is a co-atomic lattice.*

A  $q$ -PMD is a  $q$ -matroid such that all flats of the same rank have the same dimension.

Lemma

*$q$ -Steiner systems are  $q$ -PMDs.*

Fact: finding  $q$ -Steiner systems is hard. Maybe  $q$ -matroids help?

Conjecture (J. & Torielli, 2017)

All  $q$ -matroids come from rank metric codes.

That means: a  $q$ -matroid over  $E = \mathbb{F}_q^n$  of rank  $k$  can be represented by a  $k \times n$  matrix over a suitably large extension field  $\mathbb{F}_{q^m}$ .

To do list:

- ▶ Fix details.
- ▶ Do  $q$ -PMDs give us subspace designs?
- ▶ Do other results on PMDs have a  $q$ -analogue? (Deza, 1992)
- ▶ Residual/derived design vs deletion/contraction in  $q$ -matroid.
- ▶ Relation between the representation matrix and the automorphisms of a design?
- ▶ Find a representation of the  $\mathcal{S}_2(2, 3, 13)$   $q$ -Steiner system.
- ▶ Wishful thinking: what about the  $q$ -analogue of the Fano plane ... ?

Help is welcome!